



Limitations of Regulated Deficit Irrigation

Other topics in this Viti-Notes series include:

- An introduction to Regulated Deficit Irrigation
- Scheduling Regulated Deficit Irrigation
- Varietal responses to Regulated Deficit Irrigation

The ability to implement deficit irrigation strategy is influenced by climate and soil type. Most difficulties in practising Regulated Deficit Irrigation (RDI) occur where soil is difficult to dry out either because it is a type which retains moisture or it is in a region where late spring rainfall keeps it moist. Soil must dry out enough to induce a deficit response at the required growth stage and take in moisture at an adequate rate to relieve the deficit when irrigation is resumed. These are influenced by the rates of infiltration and evapotranspiration, and the total water available to vine roots. The most significant limitations to a successful RDI program are presented below.

Climatic conditions

High winter and spring rainfall and rainfall events through the growing season make it difficult to dry the soil and produce a deficit response in the vines. Regions that start the growing season with full soil moisture profiles may find difficulties drying the subsoils enough to get to a deficit situation by fruit set, especially if spring rainfall is plentiful. Managing planted mid rows to utilise excess water may help in these situations although care needs to be taken to ensure enough soil moisture is available and irrigation amounts are not increased as a result. Where rainfall patterns do not make RDI possible there may still be scope through careful timing of irrigation to exercise some control over canopy vigour.

High early summer temperatures can mean special care needs to be exercised when applying a deficit program. Experience in vineyards in hot climates indicate that RDI can be applied successfully when vines are irrigated sufficiently before flowering, particularly where soils and irrigation systems allow quick infiltration of water immediately at the conclusion of the RDI period.

Different soil types

Lighter soil types (sands, sandy loams) are the easiest soils with which to implement deficit irrigation practices as the soil moisture levels are able to be controlled quite easily (depending on climate of course). These soils generally dry out relatively quickly and also re-wet quickly. Control of soil moisture in these situations can be very precise, except where the soil profile and root extension is very deep.

Clays are more difficult to manage but with practice and experience can successfully deliver desired responses in the vine. These soils generally take longer to dry out at the beginning of the season and are then more difficult to re-wet once dry. Careful management of moisture levels in the active root zone ensures that adequate soil moisture is available all through the growing season.

It is extremely difficult to manage RDI programs in heavy soils like black cracking clays as they store large amounts of water and crack when they do dry out, making re-wetting a slow process. Proceed with caution if applying deficit irrigation on these soil types. Other practices (such as mulching) may need to be implemented to successfully manage the system. It is advisable in these situations that a very small section of the vineyard is trialled with the deficit practice until confidence in management is gained.

Duplex soils (sandy loams or loams over clay subsoils) present specific challenges. Generally the top soil acts like the lighter soils described above and the sub soil like the heavier soils, with the added difficulty of transient water tables forming over the clay subsoil if it is impermeable, or if the application rate is too high. Management of duplex soils depends on knowing vine root distribution and gaining experience with the drying and wetting characteristics of the soil.

Further information

Water Management for Grapevine Production: Research to Practice¹ is a training program whose delivery can be fine-tuned to suit each region.

- Enquiries to Peter Mansfield on (08) 8222 9255 or visit www.crcv.com.au for more information.

Numerous articles about Regulated Deficit Irrigation and other water management techniques have been published in various issues of The Australian and New Zealand Grapegrower and Winemaker. Visit www.grapeandwine.com.au/ for details.

Visit the web site at www.crcv.com.au/viticare/vitinotes/ for updates and more Vitinote titles.

Product or service information is provided to inform the viticulture industry about available resources, and should not be interpreted as an endorsement.

©2004 Cooperative Research Centre for Viticulture

THE CRC for Viticulture is a joint venture between the following core participants, working with a wide range of supporting partners.



AWRI



Know-how for Horticulture™



Australian Government
Grape and Wine Research and
Development Corporation

